

References

- Aleksidze M. A.* The solution of some fundamental problems of gravimetry. — Tbilisi: Metsniereba, 1985. — 412 p. (in Russian).
- Andó B., Carbone D.* A test on a neuro-fuzzy algorithm used to reduce continuous gravity records for the effect of meteorological parameters // *Phys. Earth Planet. Int.* — 2004. — **142**. — P. 37—47.
- Bolotnova L. A.* Eco-geological study of the state of geological environment in urban areas: geophysical aspects / V. V. Filatov, L. A. Bolotnova // *IX Geophys. readings after V. V. Fedynskiy*, 1—3 March 2007: Abstr. proceedings. — Moscow, 2007. — P. 43—44 (in Russian).
- Bychkov S. G.* On the calculation of gravity anomaly in the Bouguer reduction // *IX Geophys. readings after V. V. Fedynskiy*, 1—3 March 2007: Abstr. proceedings. — Moscow, 2007. — P. 73—77.
- Chorna O. A.* Investigation of inverse problems of the logarithmic potential theory for bodies resembling the given ones: Thesis for a cand. degree on phys.-mat. sci. / National Academy of Sciences of Ukraine. — Kiev, 1999. — 26 p. (in Russian).
- Dubovenko Yu. I.* Restoration of the contact boundary in layered medium // *Geophys. J.* — 2002. — **24**, № 6. — P. 36—41 (in Ukrainian).
- Dvulit P. D.* Methods of accounting of geophysical factors on the variation of the gravity field of the Earth. — F. Dr. theses in Engineer. Sci: 05.24.01. Lviv Polytechnical. — Lvov, 1999. — 225 p. (in Ukrainian).
- Regularization algorithms and a priori information* / Eds. A. N. Tikhonov, A. V. Goncharsky, V. V. Stepanov, A. G. Yagola. — Moscow: Science, 1983. — 200 p.
- Sobakar G. T.* Quasiperiodic variations of the gravity field of the Earth, their nature and applied scientific value // *Geophys. Proceedings AS USSR*. — 1972. — **46**. — P. 31—42 (in Russian).
- Starostenko V. I., Legostaeva O. V., Makarenko I. B., Pavlyuk E. V., Sharypanov V. M.* On the automated input into a computer the images of the geological and geophysical maps with gaps of 1st kind and the interactive visualization of 3-D geophysical models and their fields // *Geophys. J.* — 2004. — **26**, №1. — P. 3—13 (in Russian).
- Yurgin O. V.* High-precision gravity prospecting for measurement of gravitational effects of shallow origin: Thesis for a cand. degree on engineer. sci. — Perm, 2006. — 26 p. (in Russian).
- Yurkina M. I.* Definition of measurements of the gravity field and the vertical crustal movements by the repeated gravimetric and levelling observations // *Geodesy and Cartography*. — 1978. — № 4. — P. 30—35 (in Russian).

Shallow coseismic slip deficit due to large (M7) strike-slip earthquakes

© *Y. Fialko, 2010*

Institute of Geophysics and Planetary Physics, Scripps Institution of Oceanography,
University of California San Diego, San Diego, USA
yfialko@ucsd.edu

Inversions of space geodetic data (in particular, Interferometric Synthetic Aperture Radar and Global Positioning System) from several large (moment magnitude ~7) strike-slip earthquakes indicate that coseismic slip in the middle of the seismogenic layer (at depth of 4—5 km) is systematically larger than

slip at the Earth's surface. Fig. 1 shows an example of slip inversion from the April 4, 2010, M7.2 El Mayor (Mexico) earthquake, and Fig. 2 shows a compilation of slip inversions from several well-documented events [Fialko et al., 2005], including our recent results for the El Mayor earthquake.




Fig. 1. Coseismic slip model of the El Mayor earthquake derived from inversion of InSAR and GPS Figure data. Colors denote the slip magnitude and arrows denote the sense of slip on the west side of the fault. Black circles denote precisely relocated hypocenters of aftershocks from the time period of 2 months following the mainshock (courtesy of Egill Hauksson).




Fig. 2. Along-fault averaged distribution of slip for several large strike-slip earthquakes well constrained by the space geodetic data [Fialko et al., 2005]. The slip distribution from the El Mayor earthquake (magenta curve) follows the general pattern, with slip maximum in the middle of the seismogenic layer and shallow coseismic slip deficit.

This decrease in slip toward the surface, termed “shallow slip deficit”, appears to be consistent with the idea that the uppermost brittle layer is velocity-strengthening, as suggested by experimental data [Marone, 1998; Scholz, 1998], there remain a question of how the coseismic slip deficit is accommodated throughout the earthquake cycle [Fialko et al., 2005]. To the best of our knowledge, events included in Fig. 2 were not associated with either

shallow interseismic creep or robust shallow after-slip (in the amount sufficient to remove the coseismic slip deficit in the shallow crust) [Jacobs et al., 2002; Fialko, 2004; Fialko et al., 2005; Fielding et al., 2009]. We explore a possibility that the shallow slip deficit is associated with immature and/or infrequently slipping faults and is caused by the bulk inelastic yielding of the host rocks in the shallow part of the brittle crust.

References

Fialko Y. Evidence of fluid-filled upper crust from observations of post-seismic deformation due to the

1992 Mw 7.3 Landers earthquake // J. Geophys. Res. — 2004. — **109**. — P. B08,401.

- Fialko Y., Sandwell D., Simons M., Rosen P.* Three-dimensional deformation caused by the Bam, Iran, earthquake and the origin of shallow slip deficit // *Nature*. — 2005. — **435**. — P. 295—299.
- Fielding E. J., Lundgren P. R., Burgmann R., Funning G. J.* Shallow fault-zone dilatancy recovery after the 2003 bam earthquake in iran // *Nature*. — 2009. — **458**. — P. 64—68.
- Jacobs A., Sandwell D., Fialko Y., Sichoix L.* The 1999 (Mw 7.1) Hector Mine, California, earthquake: Near-field postseismic deformation from ERS interferometry // *Bull. Seism. Soc. Amer.* — 2002. — **92**. — P. 1433—1442.
- Marone C.* Laboratory-derived friction laws and their application to seismic faulting // *Ann. Rev. Earth Planet. Sci.* — 1998. — **26**. — P. 643—696.
- Scholz C. H.* Earthquakes and friction laws // *Nature*. — 1998. — **391**. — P. 37—42.

The onset of plate tectonics on super-Earth's using a damage rheology

© *B. Foley¹, D. Bercovici¹, W. Landuyt², 2010*

¹Yale University Department of Geology and Geophysics, New Haven CT, USA
bradford.foley@yale.edu
david.bercovici@yale.edu

²University of California San Diego, Scripps Institute of Oceanography, San Diego CA, USA
wlanduyt@ucsd.edu

Numerical simulations of mantle convection with a damage — grainsize feedback are used to develop scaling laws to predict conditions at which super-Earths would have plate tectonics. In particular, we introduce a new criterion for the onset of plate tectonics on terrestrial planets: that the viscosity of the lithosphere must be reduced to some critical value, which we assume to be the mantle viscosity. We formulate this criterion using the viscosity ratio between the pristine lithosphere and underlying mantle (μ_0/μ_1). These conditions are mapped out in regime diagrams of μ_0/μ_1 versus the damage fraction (f_a). The regime diagrams show that the transition from stagnant lid to mobile surface occurs for higher μ_0/μ_1 as f_a increases, with a power law relationship between those two variables; moreover, decreasing the healing constant (k_a) at the surface shifts the transition boundary to higher μ_0/μ_1 . A scaling law is developed assuming that the transition

between regimes occurs when damage, driven by convective stresses, reduces the viscosity in the lithosphere to a viscosity comparable to the mantle viscosity. This scaling law explains the numerical results well and can be applied to terrestrial planets. For the Earth, damage is efficient in the lithosphere, and viscosity can be reduced by 10 orders of magnitude with grains being reduced to a size on the order of a micron. When applied to super-Earth's, we find that larger planets are capable of larger viscosity reductions, but the viscosity ratio increases with planetary size at roughly the same rate. Therefore, contrary to previous results [e. g. O'Neill, Lenardic, 2007; Valencia et al., 2007], we find that the size of the planet has little effect on the convective regime that planet lies in. Factors such as surface temperature and thermal evolution may be more important in explaining the convective style of terrestrial planets.

References

- O'Neill C., Lenardic A.* Geological Consequences of Super-sized Earths // *Geophys. Res. Lett.* — 2007. — **34**. — P. 19204—19208.
- Valencia D., O'Connell R. J., Sasselov D. D.* Inevitability of Plate Tectonics on Super-Earths // *Astrophys. J.* — 2007. — **670**, № 1. — P. L45—L48.